

NEUDISIN

Nuetral Disinfectant Cleaner

Efficacy

Hospital Disinfection (at 2 ounces per gallon)

NEUDISIN is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum and 400 ppm hard water with a 10 minutes contact time against:. (Testing is performed per the AOAC UDT/ GST method (DIS/TSS-1). Sixty carriers are required on 3 separate lots, one of which must be > 60 days old against Pseudomonas aeruginosa, Salmonella enterica and Staphylococcus aureus. Killing of 59 out of 60 carriers is required (total carriers = 540).)

ORGANISM	CARRIER POPULATION	SAMPLE	# CARRIERS	# POSITIVE
Pseudomonas aeruginosa ATCC #15442	3.9 X 10 ⁶ CFU/Carrier	A	60	0/60
		(60 Days Old)		
		B	60	0/60
Salmonella enterica ATCC #10708	1.03 X 10 ⁸ CFU/Carrier	C	60	1/60
		(60 Days Old)		
		A	60	1/60
Staphylococcus aureus ATCC #6538	7.0 X 10 ⁸ CFU/Carrier	B	60	1/60
		C	60	0/60
		A	60	0/60
(60 Days Old)				
		B	60	0/60
		C	60	0/60
		A	60	0/60

Supplemental Organisms

(Testing is performed per the AOAC UDT/GST method. Ten carriers are required on 2 separate lots against each supplemental organism. Killing of 10 out of 10 carriers is required (total carriers = 20).)

ORGANISM	CARRIER POPULATION	SAMPLE	# CARRIERS	# POSITIVE
Acinetobacter baumannii ATCC 19003	5.1 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Acinetobacter Iwoffi ATCC 15309	5.7 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Bordetella bronchiseptica ATCC 10580	9.4 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Citrobacter freundii ATCC 8090	3.9 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterobacter aerogenes ATCC 13048	2.35 X 10 ⁷ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterobacter agglomerans ATCC 27155	3.9 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterobacter cloacae ATCC 13047	3.3 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterococcus faecalis ATCC 19433	6.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterococcus faecalis Vancomycin Resistant (VRE) ATCC 51299	1.3 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Enterococcus hirae ATCC 10541	1.19 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Escherichia coli ATCC 11229	1.3 X 10 ⁷ CFU/Carrier	A	10	0/10
		B	10	0/10
Escherichia coli (carbamapem resistant) CDC 87371	1.34 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Escherichia coli Tetracycline Resistant ATCC 47041	3.1 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Escherichia coli Spectrum B-Lactamase (ESBL) ATCC BAA-196	4.6 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Fusobacterium necrophorum ATCC 27852	5.8 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Klebsiella oxytoca ATCC 13182	1.07 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Klebsiella pneumoniae ATCC 13883	1.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Klebsiella pneumoniae (New Delhi metalo-beta-lactamase) CDC1000527	9.5 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Listeria monocytogenes ATCC 19117	7.7 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Micrococcus luteus ATCC 14452	1.1 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Pasturella multocida ATCC 12947	1.32 X 10 ⁷ CFU/Carrier	A	10	0/10
		B	10	0/10
Proteus vulgaris ATCC 13315	1.9 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Pseudomonas aeruginosa Tetracycline Resistant ATCC 27853	3.5 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Pseudomonas cepacia ATCC 25416	1.63 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Salmonella enterica ATCC 23564	9.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Salmonella enterica serotype pullorum ATCC 19945	7.1 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Salmonella typhi ATCC 6539	8.3 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Salmonella typhimurium ATCC 23564	1.5 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Serratia marcescens ATCC 14756	6.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Shigella flexneri ATCC 9380	1.99 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Shigella sonnei ATCC 25931	1.04 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus aureus ATCC 25923	6.6 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus aureus sub species aureus ATCC 33586	7.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus aureus Methicillin Resistant (MRSA) ATCC 33592	5.4 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus aureus Community Associated Methicillin Resistant (CA-MRSA)	6.3 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus aureus Vancomycin Intermediate Resistant (VISA) ATCC 5836	3.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus epidermidis ATCC 14990	1.56 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus epidermidis Antibiotic resistant ATCC 51625	8.6 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Staphylococcus haemolyticus ATCC 29970	9.5 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Streptococcus agalactiae ATCC 13813	5.6 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10



- Virucide*
- Mildewstat
- Fungicide

Efficacy Data

EPA Reg. No. 10324-154-72072

EPA Est. No. 1459-PA-01

Product Code: BS300-01

ORGANISM	CARRIER POPULATION	SAMPLE	# CARRIERS	# POSITIVE
Streptococcus mutans ATCC 25175	1.02 X 10 ⁹ CFU/Carrier	A	10	0/10
		B	10	0/10
Streptococcus pneumoniae Penicillin Resistant ATCC 51915	9.6 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Streptococcus pyogenes ATCC 19615	4.7 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Vibrio cholera ATCC 11623	1.0 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Yersinia enterocolitica ATCC 23715	1.2 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10

(Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.)

ORGANISM	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Chlamydia psittaci ATCC VR-125	7.25 Log ₁₀	A	≤0.5 Log ₁₀	≥6.75 Log ₁₀
		B	≤0.5 Log ₁₀	≥6.75 Log ₁₀
		C	≤0.5 Log ₁₀	≥6.75 Log ₁₀

Virucidal against (at 2 ounces per gallon)

NEUDISIN was evaluated in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces. (Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.) (3 lots and 4-Log reduction for Canada.)

ORGANISM	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Avian Influenza A (H3N2) virus (Avian Reassortant) (ATCC VR-2072)	4.75 Log ₁₀	A	≤0.5 Log ₁₀	≥4.25 Log ₁₀
		B	≤0.5 Log ₁₀	≥4.25 Log ₁₀
		C	≤0.5 Log ₁₀	≥4.25 Log ₁₀
Avian Influenza A (H5N1) virus	6.75 Log ₁₀	A	≤0.5 Log ₁₀	≥6.25 Log ₁₀
		B	≤0.5 Log ₁₀	≥6.25 Log ₁₀
Cytomegalovirus ATCC VR-538	4.5 Log ₁₀	A	≤0.5 Log ₁₀	≥4.0 Log ₁₀
		B	≤0.5 Log ₁₀	≥4.0 Log ₁₀
		C	≤0.5 Log ₁₀	≥4.0 Log ₁₀
Hepatitis B Virus	5.06 Log ₁₀	A	0.27 Log ₁₀	4.79 Log ₁₀
		B	0.41 Log ₁₀	4.79 Log ₁₀
		Confirmatory B	0.27 Log ₁₀	4.79 Log ₁₀
Hepatitis C Virus	6.21 Log ₁₀	A	0.24 Log ₁₀	5.97 Log ₁₀
		B	0.42 Log ₁₀	5.79 Log ₁₀
		Confirmatory B	0.13 Log ₁₀	5.93 Log ₁₀
Herpes Simplex Virus Type 1 ATCC VR-773	5.5 Log ₁₀	A	≤0.5 Log ₁₀	≥5.0 Log ₁₀
		B	≤0.5 Log ₁₀	≥5.0 Log ₁₀
Herpes Simplex Virus Type 2 ATCC VR-734	6.0 Log ₁₀	C	≤0.5 Log ₁₀	≥5.5 Log ₁₀
		A	≤0.5 Log ₁₀	≥5.5 Log ₁₀
		B	≤0.5 Log ₁₀	≥5.5 Log ₁₀
Human Coronavirus ATCC VR-740	5.75 Log ₁₀	C	≤0.5 Log ₁₀	≥4.5 Log ₁₀
		A	≤0.5 Log ₁₀	≥4.5 Log ₁₀
		B	≤0.5 Log ₁₀	≥4.5 Log ₁₀
Human Immunodeficiency Virus type 1 (HIV 1) HTLV-III _g	4.5 Log ₁₀	C	≤0.5 Log ₁₀	≥4.0 Log ₁₀
		A	≤1.5 Log ₁₀	≥4.25 Log ₁₀
		B	≤1.5 Log ₁₀	≥4.25 Log ₁₀
Influenza A virus ATCC VR-544	6.5 Log ₁₀	A	≤0.5 Log ₁₀	≥6.0 Log ₁₀
		B	≤0.5 Log ₁₀	≥6.0 Log ₁₀
		C	≤0.5 Log ₁₀	≥5.5 Log ₁₀
Influenza A (H1N1) virus ATCC VR-1469	5.5 Log ₁₀	A	≤0.5 Log ₁₀	≥5.0 Log ₁₀
		B	≤0.5 Log ₁₀	≥5.0 Log ₁₀
Respiratory syncytial virus ATCC VR-26	4.5 Log ₁₀	A	≤0.5 Log ₁₀	≥4.0 Log ₁₀
		B	≤0.5 Log ₁₀	≥4.0 Log ₁₀
		C	≤0.5 Log ₁₀	≥4.5 Log ₁₀
SARS Associated Coronavirus	6.23 Log ₁₀	A	≤3.5 Log ₁₀	≥3.0 Log ₁₀
		B	≤3.5 Log ₁₀	≥3.0 Log ₁₀

Fungicidal against (at 2 ounces per gallon)

NEUDISIN was evaluated in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time and found to be effective against the following fungi on hard nonporous environmental surfaces. (Testing is performed per the AOAC fungicidal method (DIS/TSS-6). Two separate lots are tested against Trichophyton mentagrophytes in a suspension test. Killing of all fungal spores in 10 minutes is required.)

ORGANISM	CARRIER POPULATION	SAMPLE	# CARRIERS	# POSITIVE
Candida albicans ATCC #10231	1.57 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10
Trichophyton mentagrophytes ATCC #9533	1.10 X 10 ⁸ CFU/Carrier	A	10	0/10
		B	10	0/10

Mold and Mildew Control (at 2 ounces per gallon)

Use NEUDISIN to control the growth of mold and mildew and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air dry. Repeat application weekly or when growth or odor reappears.

ORGANISM	TILE NUMBER	UNTREATED AFTER 7 DAYS	SAMPLE A AFTER 7 DAYS	SAMPLE B AFTER 7 DAYS
Aspergillus niger ATCC #16404	1	Growth 90%	No Growth 0%	No Growth 0%
	2	Growth 70%	No Growth 0%	No Growth 0%
	3	Growth 90%	No Growth 0%	No Growth 0%